

The effectiveness of increasing the volume of training load on the period of overcompensation and its effect on "breaking" in the training session after stopping for young boxers in some physical and functional variables

* DR /Ahmed Abdel Hamid Ali Zahran

Introduction and research problem:

It is known that the individual not only has the ability to carry out training loads, adapt to them, but also compensate for the excess by reaching a better level after the recovery phase according to the Law of increasing training load (17-10:12)

During the past few years, the training volumes for elite and international players have doubled, reaching between 1400-1000 training hours, and the volume of training for competitions has also increased in the training year(80-22:10) .

Accordingly, increasing the size of the training load as one of the components of the pregnancy during the training periods that range from 1-3 weeks contributes to early reaching the required physical level, changing the components of the body, and enhancing achievement. ((50-30:22)

There is also a tendency that increasing the volume of the training

load for more than two weeks during the training periods produces manifestations of excessive training load, which causes stress and a decrease in the level of achievement because it does not allow recovery to be restored, and they suggest returning to the normal training volume gradually for a period of 0-2 weeks(79-74:21) .

In order to benefit from the quantitative and qualitative training loads, the contents of the training load must be directed to achieve specific goals with gradation and diversity in the implementation of the training load from multi-directional to bi-directional and then to a single or single direction so that the player can adapt and overcompensate by reaching a better level after the recovery period according to the law of return or yield (17:14) .

And since the optimal training case "sports format" passes through three stages, the first stage is building,

*Lecturer, Department of Theories and Applications of Battles and Water Sports, Faculty of Physical Education, Sadat City University

shaping, and developing the training case, which is matched by the training course with three preparatory "preparatory" periods represented by general preparation and special preparation, and they constitute about 50% of the time. The training course and the preparation period for competitions, which constitutes 20% of the training course time. The second stage is the stage of stabilizing and maintaining the optimal training situation, and it corresponds to the competitions period, which constitutes about 10% of the training course time, and then the third stage, which is the temporary decline of the training state, and Represented by the transitional "recovery" period of the training course and constitute about 10% of the time of the training course. (2009-2008:26)

Given that one of the most important obstacles facing us in reaching the optimal training situation, "training format", and achieving the best possible level of achievement during its second phase is the presence of relatively long pauses as a result of the exams for the first and second semesters, which precede the competitions period in the training course, which is represented by In the "winter" shield competitions and the

"summer" cup, which are organized by the Egyptian Athletics Federation and its branches in the regions, which impedes the gradual, sequential, and regular progression of the contents of the training load, and training hours over the timeframe necessary to download and reach the optimal training state and achieve The best possible level of achievement in the allotted time (2007-2006:3).

Accordingly, fitness and achievement improve as a result of the correct relationships between the training load and adapting to it according to the Law of Reversibility. Therefore, it is necessary to be aware of the problems that obstruct this, and to solve them creatively so that we can take decisions and actions that achieve the goals set in their specific times. (2007:11)

Through the researcher's work as a boxing coach, and during the players' pause and interruption of training in order to study and exams, especially starting from the month of June, he noticed the low level of performance of young boxers, which may be due to their lack of some physical and functional qualities. Undoubtedly, there is more than one way to achieve this. But the question

comes, what are the best ways to overcome these problems with a great deal of success to reach the best possible level of achievement during the competition period, and accordingly the research problem crystallized in the need to identify how to “double” the size of the training load during the period following the period of cessation of training, and Until the competitions period, and its effect on some physical and functional variables.

research aims--:

1- Develop an appropriate planning for the formation of the contents of the training load cycle with an increase in the size of the training load that can be used during the time available for download after the breaks and until the date of the main competitions.

2-To identify the effect of using an increase in the size of the training load on some physical and functional variables of young boxers

Research hypotheses :-

1- There are statistically significant differences between the tribal and remote measurements in favor of the post measurement in the physical variables of young boxers

2- There are statistically significant differences between the pre and post measurements in favor of the post measurement in the functional variables of young boxers

Search terms:-

1- Increasing the training load: It is to increase the levels of loading until reaching the outer limit of the player's ability (affective fatigue), which leads to increased adaptation and overcompensation to a more advanced level. (١٢-٣٢:٧)

Search procedures :-

The experimental method was used using the experimental design of two groups, one experimental and the other control.

The research sample

The research sample was chosen by the intentional method of youth boxing players from above 17: less than 19 years old from clubs and centers (Al-Nujoum Sports Club - Al-Salam Club - Sadat City Youth Center) in Al-Menoufia region. The number of the research sample members was ١٧ young boxers, including ٦ boxers who were subjected to the exploratory study, ١١ boxers for the basic experience as shown in the table

Table (1)
Classification of the total population of the research sample

| م | sports institutions | the statement | | |
|---|-------------------------|--------------------|--------------|-------------|
| | | Experimental group | Experimental | group Total |
| ١ | Al-Nujoom Sports Club | ٣ | ٢ | ٥ |
| ٢ | Peace Club | ٥ | ٢ | ٨ |
| ٣ | Sadat City Youth Center | ٣ | ٢ | ٥ |
| ٥ | the total | ١١ | ٦ | ١٧ |

Reasons for choosing the research sample

- Regular players in terms of commitment and training dates.
- The willingness of all players to attend training to participate in the research group.
- Convergence of time, training, physical and technical capabilities of

the sample, as well as the educational and social level.

The homogeneity of the research sample

The researcher performed homogeneity on the basic research sample before applying the training program in growth rates (age – height – weight – training age) and Table No. illustrates this (2)

Table (2)
The arithmetic mean, median, standard deviation, and skewness coefficient for an individual .The research sample in the variables (age, height, weight, training time) n = 17

| | Variables | | mean | median | deviation | skew |
|---|--------------|------------|--------|--------|-----------|--------|
| ١ | Age | year | ١٧,٨٧ | ١٨ | ٠,٦٤٠ | ١,٤٨- |
| ٢ | Height | Centimeter | ١٧٥,٣٦ | ١٧٥,٥ | ٥,٣٩٠ | ٠,٠٨٠- |
| ٣ | Weight | Kg | ٦٦,٢٧ | ٦٤ | ١٠,٦٦ | ٠,٦٤٠ |
| ٤ | training age | Month | ٣,٥٤ | ٣ | ١,٢٦ | ١,٣٠ |

It is clear from Table No. ٢ that all the torsion coefficients of the research sample ranged between (-١.٤٨٠, ١.٣٠) and that these values were

limited to between +٣, which confirms the homogeneity of the research sample in the variables (age – height – weight – training age).

Data collection methods

The researcher used a variety of methods to collect data in line with the nature of the research and the data to be obtained, which are as follows:

The player's personal data registration form and physical and physiological tests.

Tests used:

By reviewing the scientific references and previous studies, tests that measure physical characteristics and physiological variables were extracted Selection of assistants:

- Six assistants were selected from the graduates of the Faculty of Physical Education, who were prepared for this by the researcher's knowledge.
- Survey Studies

The exploratory study was conducted from ۲۰۲۱/۹/۲ to ۲۰۲۱/۱۸/۹ on a representative sample of the indigenous community and from outside the basic research sample, which consisted of ۱۲ boxers, including ۶ distinguished youth boxers from

among the heroes of the republic, ۶ undistinguished youth boxers of the same age group and targeted the following Finding validity and reliability coefficients for physical and physiological tests.

The results of the second survey:

- The stability coefficient of physical and physiological tests was found by applying the test and reapplying it with a time difference of ۳۰ minutes as shown in Table No. ۱۰. The stability coefficient was found for it over three days, where the researcher divided the boxers into three groups according to weight (first day light weights – today The second day (medium weights, then the third day, heavy weights).

The validity coefficient (discrimination validity) was found for the physical and physiological tests by using the second measure of the reliability coefficient and comparing it with the results of ۶ distinguished young boxers of the same age group. Table No. ۳ shows the validity coefficient

(۳) Table

The reliability coefficient of the tests in question n = n2 = 6

| ردیف | Test name | First application | | Second application | | application Correlation coefficient |
|------|---|-------------------|-----------|--------------------|-----------|-------------------------------------|
| | | mean | deviation | mean | deviation | |
| ۱ | Throw a 3 kg medicine ball to the farthest distance with the right hand | ۷,۵ | ۰,۹۳۹ | ۷,۳۷ | ۰,۶۸۱ | *۰,۶۶۴ |
| ۲ | Throw a 3 kg medicine ball to the farthest distance with the right hand | ۶,۹۴ | ۱,۶۵۴ | ۶,۴۸۴ | ۱,۳۶۸ | *۰,۸۲۸ |

| | | | | | | |
|---|--|--------|--------|--------|--------|--------|
| ٣ | Left and right straight shot on the punching bag within (20 seconds) | ٤٤,٣ | ٥,٢٩٢ | ٤٢,٩ | ٤,٢٥٤ | *٠,٨٧٥ |
| ٤ | Straight left and right shot on the punching bag within (1 s) | ٨٩,٣ | ٨,٩٢٠ | ٨٩,١ | ١٠,١١٥ | *٠,٨٥٥ |
| ٥ | Moving forward and backward between the flags with straight punches left and right to the head | ١١,٣٤ | ١,١٢٧ | ١١,٤٩ | ١,٠٦٣ | *٠,٨٢٠ |
| ٦ | Maximum oxygen consumption | ٣,٢٤٦ | ٠,١١٥ | ٣,١٩٠ | ٠,١٣١ | *٠,٧١٠ |
| ٧ | physical efficiency | ٧٧,١١ | ٣,٤٨٤ | ٧٩,١٧٢ | ٦,٠٧٥ | *٠,٨٦١ |
| ٨ | anaerobic capacity | ٣٨٩,١٤ | ٤٤,٠٣٤ | ٣٩٠,٥٢ | ٤٣,٣٥ | *٠,٨٩٠ |

(t) tabular value at the level of 0.05 = 0.602

It is clear from Table No. 3 that the correlation coefficients for the tests were limited to (0.644, 0.890) and that

all correlation coefficients are significant at the 0.05 level, which confirms the stability of the tests.

table(4)

The differentiation validity coefficient of the tests under study n1 = n2 = 6

| ٨ | Test name | Distinguished group | | A less group | | distinguishes | teams value |
|---|--|---------------------|-----------|--------------|-----------|---------------|-------------|
| | | Mean | deviation | mean | deviation | | |
| ١ | Throw a 3 kg medicine ball to the farthest distance with the right hand | ٩,٦٥ | ١,٧٦٦ | ٧,٣٧٣ | ٠,٦٨١ | ٢,٢٨٠ | *٣,٦١٠ |
| ٢ | Throw a 3 kg medicine ball to the farthest distance with the right hand | ٨,٢٥ | ١,٣٨٨ | ٦,٤٨٤ | ١,٣٦٨ | ١,٧٦٦ | *٢,٧٢٠ |
| ٣ | Left and right straight shot on the punching bag within (20 seconds) | ٦٥ | ١٢,٦٨ | ٤٢,٩ | ٤,٢٥٤ | ٢٢,١ | *٤,٩٦٠ |
| ٤ | Straight left and right shot on the punching bag within (1 s) | ٩٩,٩ | ٧,٩٠٨ | ٨٩,١ | ١٠,١١٥ | ١٠,٨ | *٢,٥٢ |
| ٥ | Moving forward and backward between the flags with straight punches left and right to the head | ١٠,٠٧ | ٠,٣٦٦ | ١١,٤٩ | ١,٠٦٣ | ١,٤٢ | *٣,٧٠ |
| ٦ | Maximum oxygen consumption | ٤,٠٧٤ | ٠,٣٧٨ | ٣,١٩ | ٠,١٣١ | ٠,٨٨٤ | *٦,٦١٠ |
| ٧ | physical efficiency | ٩٣,٨١٢ | ٦,٨٤١ | ٧٩,١٧٢ | ٦,٠٧٥ | ١٤,٦٤ | *٤,٨٠ |

T-table value at the level of 0.05 = 2.101

It is clear from Table No.4 that the validity coefficient is highly significant between the privileged and undistinguished group, where the degree of honesty is between (2.52, 6.61) and that all of them are significant at the 0.05 level, which confirms the validity of these tests in what they were designed for, and that it can differentiate between distinguished and non-distinguished boxers. Distinguished in the same age group.

Basic study :

Tribal measurements:

Tribal measurements were carried out for the members of the research sample (the experimental group) in the period from 2021/24/9 to 2021/29/9 at the Boxing Hall at the Faculty of Physical Education in Sadat City. These measurements took (5) days and included: (age – height – weight – age Training – physical tests – physiological tests)

Executing the experiment:

The training program was implemented on the experimental group for a period of one and a half months, from 2021/10/1 to 2021/10/11, with 8 training units per week.

training program:

The aim of the program:

The training aims to increase the size of the training load on the period of overcompensation and its effect on the “fracture” in the training session after stopping for young boxers in some physical and functional variables.

Program foundations:

The researcher laid the foundations of the program after reviewing the scientific references specialized in boxing, training science and previous studies as follows:

–The duration of the training program is 6 weeks.

–The number of weekly training units has been set at 8 training units.

–The time of the training unit was limited to the week with medium load between (100-120s), in the week with high load by (90- 120s), and in the week with maximum load by (110- 130s), provided that the inter-rest periods are large, which leads to an increase training unit time.

The load degrees were distributed over the training weeks during the training program, provided that the average load is between (50-70%), the high load is between (70-80%) and the maximum load is between (80: to the limits of the player's amount 100%).

-The researcher took into account the distribution of the degree of pregnancy between weeks and levels of training intensity in a ratio of (1:1) in line with the requirements of competition in boxing and what it requires from the necessity of performance in tournaments over the

course of the championship days in succession.

The researcher relied on calculating pulse rates to determine the intensity of the training load, through the following physiological trends

Table (5)

Pregnancy types and training trends used in the research

| Pregnancy types | Average | High | Maximum |
|---|------------------------|------------------------|------------------------|
| Measurements | z/min 100 - 130 | z/min 170 - 190 | z/min 200 - 170 |
| heart rate | to 2 degrees 1 | to 4 degrees 2 | ie 10 degrees 3 |
| What is the equivalent in ?degrees | from 30 - 60% | from 60 - 90% | %100 - 80 |

Pulse rates can be calculated using the maximum heart rate reserve (HRR) method, which is the difference between the athlete's maximum heart rate during exercise and his resting pulse rate. This can be calculated as follows:

-Maximum heart rate: $220 - (\text{average age})$.

Reserve max heart rate = max resting heart rate - max resting heart rate (reserve maximum pulse rate)
 $HRR = \text{max HR} - \text{rest HR}$

Required intensity \times pulse reserve

Average heart rate = + resting heart rate 100

The gradual upgrading of the solution level was taken into account through the systematic change of its components.

This does not mean increasing the training load daily, and in general, regulating the times of the loads

The training courses and their distribution on the training weeks and the appropriate degrees of pregnancy during the duration of the program according to Table No. 15

Table (6)
Training times and pregnancy degrees during the weeks of the program

| the week | | ١ | ٢ | ٣ | ٤ | ٥ | ٦ | ٧ | the total | pregnancy degree |
|------------------------------|----------------------|-----|-----|-----|-----|-----|-----|-----|-----------|------------------|
| the first | by level | ٩٠ | ١٢٠ | ٩٠ | ١٢٠ | ١٠٥ | ١٣٥ | ٩٠ | 750 | Average |
| Second | for the first %١٥ + | ١٠٥ | ١١٥ | ١٠٥ | ١١٥ | ١٢٠ | ١٢٥ | ١٠٠ | 785 | high |
| Third | for the second %١٥ + | ٧٥ | ٧٥ | ٧٥ | ٧٥ | ٧٥ | ١٠٥ | ٧٥ | 555 | Maximum |
| the fourth | for the third %١٥ + | ١٠٥ | ١١٥ | ١٠٥ | ١١٥ | ١٢٠ | ١٢٥ | ١٠٠ | 785 | Maximum |
| Fifth | High load for boxing | ١٢٠ | ٩٠ | ١٢٠ | ١٠٥ | ١٣٥ | ١٢٠ | ١٢٠ | 810 | high |
| VI | like the fifth | ٧٥ | ٩٠ | ٩٠ | ٩٠ | ١٠٥ | ٩٠ | ٧٥ | 615 | high |
| Total time in minutes | | | | | | | | | 4300 | |
| Total time in hours | | | | | | | | | 71.66 | |

I forced the fractions into a whole number in favor of the training program

The researcher took into account the principles and foundations of developing physical attributes, the most important of which is the gradual development.

The form of performance in the exercises is similar to the nature of performance in the sport of boxing.

-The researcher took into account the individual differences between the players, as he divided them into three groups according to weight (light - medium - heavy) during the development of physical characteristics.

Program content:

The training program includes a set of exercises directed to the development of the physical characteristics selected according to the reference survey and expert opinion poll during general and private physical preparation and preparation for matches, which were included in the training stages for the experimental group.

The program included training on straight punches in the head and linking them in the form of two and three sets, as well as some defenses for straight punches as well as the counter-attack of those punches. The program also included some tactical aspects

of the ring art, with the aim of improving all the selected research variables (physical - physiological).

Program setting steps:

1-The researcher conducted a reference survey of some scientific references specialized in the sport of boxing and the science of sports training, in addition to the related studies, in order to determine the physical characteristics of the boxing players.

2- The researcher designed an opinion poll form to determine the relative importance of the composite physical characteristics for the age group under study.

3- The researcher chose the general and private physical preparation exercises that were included in the training program as a training content and applied to the boxers in the research sample.

4- The time period of the program was divided into three phases:

-The first phase of the program lasts for (1) weeks: it aims to prepare the general and push the pregnancy before the refraction

The second phase of the program lasts for two weeks:

This stage aims to develop and develop the special physical qualities and mastery of the boxer's motor skills, which helps to improve the skillful and tactical performance, which determines the level of the boxer's training status and the level of his progress. At the end of this stage, the boxer usually reaches the sports form.

The third phase of the program lasts for 4 weeks:

It aims to maintain the level in the pre-match period.

- The researcher distributed the total training times and degrees of pregnancy during the weeks of the program and the percentages on the three stages of preparation (general physical - special - skill and planning) during the preparation period for the training program according to what was mentioned in the scientific references and previous studies related to boxing training programs and table No. (16) and figure Figure (1, 2) shows the distribution of pregnancy during the weeks of preparation and the percentages on the three stages of preparation (physical - general - special - skill - planning) during the preparation period for the program.

Presentation and discussion of the results:

Table (7)
The significance of the differences between the mean of the tribal and dimensional measurements of the experimental group In physical and physiological tests n = 11

| P | Variables | Test name | Median tribal | Median dimensional | Positive ranks | | Negative ranks | | Value (z) |
|----|-------------------------|--|---------------|--------------------|----------------|------------|----------------|------------|-----------|
| | | | | | Average Rank | Total Rank | Average Rank | Total Rank | |
| 1 | The power of speed | Throwing a 3 kg medicine ball with the right hand to the farthest distance | 8,204 | 11,0 | 6,00 | 66,00 | - | - | 2,946 |
| | | Throwing a 3 kg medicine ball with the left hand to the farthest distance | 6,909 | 9,36 | 6,00 | 66,00 | - | - | 2,900 |
| 2 | bearing speed | Left and right straight shot on punching bag (20 seconds) | 44,90 | 96,00 | 6,00 | 66,00 | - | - | 2,943 |
| 3 | endurance force | Shooting at the punching bag for (1 min) | 87,00 | 168,18 | 6,00 | 66,00 | - | - | 2,936 |
| 4 | performance endurance | Punching the left and right straight punches to the head on different punching devices | 2,307 | 1,110 | - | - | 6,00 | 66,00 | 2,936 |
| 5 | Compatibility | Throwing straight punches to the left and right on the numbered circles on the boxing sign | 14,782 | 8,012 | - | - | 6,00 | 66,00 | 2,934 |
| 6 | agility | Moving forward and backward between the flags while hitting the left and right punches to the head | 12,804 | 9,439 | - | - | 6,00 | 66,00 | 2,934 |
| 7 | physiological variables | Maximum oxygen consumption | 3,407 | 4,191 | 6,00 | 66,00 | - | - | 2,934 |
| 8 | | physical aptitude | 79,132 | 114,66 | 6,00 | 66,00 | - | - | 2,934 |
| 9 | | anaerobic capacity | 428,4 | 524,94 | 6,00 | 66,00 | - | - | 2,934 |
| 10 | | Anaerobic capacity | 378,0 | 520,6 | 6,00 | 66,00 | - | - | 2,934 |
| 11 | | lactic acid | 9,104 | 4,940 | 1,00 | 1,00 | 6,00 | 60,00 | 2,847 |

(y) tabular value at the level of 0.05 = + 1.96

It is clear from Table No. 7 that there are statistically significant differences between the mean of the tribal and dimensional measurements of the

experimental group in favor of the post-measurement at the level (0,00) in the tests of physical traits and physiological variables

Table (8)
The percentage of improvement in the experimental group in the traits tests
physical and physiological variables

| م | physical character | test name | measur | measur | of the | Improv |
|----|-------------------------|--|--------|--------|--------|---------|
| | | | ement | ement | | |
| | | | س | س | averag | percent |
| | | | | | es | % age |
| 1 | The power of speed | Throwing a 3 kg medicine ball with the right hand to the farthest distance | ٨,٢٥٤ | ١١,٥ | ٣,٢٤٦ | ٣٩,٣ |
| | | Throwing a 3 kg medicine ball with the left hand to the farthest distance | ٦,٩٠٩ | ٩,٣٦ | ٢,٤٥١ | ٣٥,٥ |
| ٢ | bearing speed | Left and right straight shot on punching bag (20 seconds) | ٤٤,٩٠ | ٩٦,٠٠ | ٥١,١ | ١١٣,٧ |
| ٣ | endurance force | Shooting at the punching bag for (1 min) | ٨٧,٠٠ | ١٦٨,١٨ | ٨١,١٨ | ٩٣,٣ |
| ٤ | performance endurance | Punching the left and right straight punches to the head on different punching devices | ٢,٣٥٧ | ١,١١٠ | ١,٢٤٧ | ٥٢,٩ |
| ٥ | Compatibility | Throwing straight punches to the left and right on the numbered circles on the boxing sign | ١٤,٧٨٢ | ٨,٥١٢ | ٦,٢٧ | ٤٢,٤ |
| ٦ | agility | Moving forward and backward between the flags while hitting the left and right punches to the head | ١٢,٨٠٤ | ٩,٤٣٩ | ٣,٣٦٥ | ٢٦,٣ |
| ٧ | physiological variables | Maximum oxygen consumption | ٣,٤٠٧ | ٤,١٩١ | ٠,٧٨٤ | ٢٣,٠٠ |
| ٨ | | physical efficiency | ٧٩,١٣٢ | ١١٤,٦٦ | ٣٥,٥٢٨ | ٤٤,٩ |
| ٩ | | anaerobic capacity | ٤٢٨,٤ | ٥٢٤,٩٤ | ٩٦,٥٤ | ٢٢,٥ |
| ١٠ | | Anaerobic capacity | ٣٧٨,٥ | ٥٢٠,٠٦ | ١٤١,٥٦ | ٣٧,٤ |
| ١١ | | lactic acid | ٩,١٥٤ | ٤,٩٤٥ | ٤,٢٠٩ | ٥٤,٦ |

It is evident from Table ٨ that the highest percentage of improvement was for the velocity endurance component of the test (straight left and right shooting on the punching bag during “ ٢٠ seconds”) and its amount

(٪١١٣,٧), and the lowest percentage of improvement was for the anaerobic capacity variable and its amount (٪٢٢,٥)

It is clear from Table No. (٣١), (٣٢) and Figure (١١), (١٢) that there are

statistically significant differences at the level ($\alpha = 0.05$) between the tribal and remote measurements of the experimental group in favor of the post-measurement in the choices of the composite physical characteristics under study, where the value of (Z) favoritism between ($2.934, 2.950$) while the tabular (Z) was ($+1.96$), as well as the percentage of increase ranging between ($26.3, 113.7$), which indicates the progress of the level of the experimental group players in the complex physical characteristics under study, and the researcher may refer This progress is due to the effect of the training program, which depends on regular training, as well as the number of training units for this group, which amounted to (4) training units per week, and the continuation of the training period.

It is clear from Table (Δ) that there is a discrepancy in the percentage of improvement from one component to another, so we find that the highest percentage of improvement was for speed endurance and was (113.7), followed by endurance of force and it was (93.3), then performance tolerance by (52.9), followed by compatibility with (42.4), then the speed advantage of the "right arm" by

(39.3), then the left arm by (35.5), and the lowest percentage of improvement for agility was (26.3)

The researcher may attribute the progression of the experimental group's player level in the composite physical attributes in question to the effect of the training program that contained different training groups aimed at developing the compound physical attributes, whether it was individual exercises based on body weight or marital based on the weight of a colleague or exercises using medical balls and exercises Weights, as well as training on devices and tools that are similar to playing situations, taking into account the individual differences of the loads, as well as the selection of muscle exercises according to muscle contraction, which led to an increase in the strength of working muscles, which contributed significantly to the improvement of the complex physical characteristics of the study sample.

The researcher attributed the result of the significance of the differences in the choice of endurance speed that modern boxing is the use of the computer depends heavily on speed more than any other element and this is consistent with what was indicated by Ismail Hamed et al. Second, third and

fourth in the same area and in sight, as well as the superiority of the Cuban players in the element of endurance of speed. (٧:١٣٤)

The researcher also attributed the high level of speed endurance due to the increase in the efficiency of the player's devices through regular training and the inclusion of the training program on playing exercises on light devices such as (dancing ball, hanging ball, bouncing ball, speed ball) as well as individual exercises to develop speed endurance during the preparation stage. The general and private physical, and confirmed by Thomason and others, W et al ١٩٩٢AD, Abdel Fattah Khader ١٩٩٦AD, that light boxing exercises affect the vital organs of the boxer in a positive way in relation to the direction of physiological work. ((٨٣:٤٩ (٣١:٤٧٢٤٦)

The results of this study agree with the results of the studies of Sameh Abdel Raouf ١٩٩٢AD (١٩) Stoletki ١٩٨٩AD (٨١) Ahmed Amin ١٩٩٥AD (٤) Arafa al-Sayed ١٩٩٧AD (٣٣) Ahmed Salah ٢٠٠٠AD (٥) Kamal Abdel-Jaber ٢٠٠٠AD (٣٦), Mustafa Nasr ٢٠٠٢AD (٥١) Yassin Ahmed ٢٠٠٢AD (٥٧), Saeed Jamil ٢٠٠٤AD (٢٢), Mustafa Amin ٢٠٠٦AD (٥٠) that the proposed training programs as well

as weight programs using forms of resistance less than the maximum, as well as special exercises, whether using boxing devices and tools or without tools leads To develop the special physical qualities of boxers.

It also appears from Table (٧) that there are statistically significant differences at the level (٠,٠٥) between the pre- and post-measurement of the experimental group in favor of the post-measurement in the physiological variables under research, where the value of (Z) favoritism was limited to (٢,٨٤٧, ٢,٩٣٤) while (Z) was tabular (+١,٩٦), as well as the percentage increase ranging between ,٪٢٢,٥) .(٪٥٤,٦

It is clear from Table No. (٧) that there is a discrepancy in the percentage of improvement from one variable to another, so we find that the highest percentage of improvement was for the lactic acid variable by (٪٥٤,٦), followed by the physical efficiency by (٪٤٤,٩), then the anaerobic capacity by (٪٣٧,٤), followed by the maximum For oxygen consumption (٪٢٣,٠), the lowest percentage of variable improvement was anaerobic capacity by.(٪٢٢,٥)

The researcher attributed the improvement in the physiological

variables under study to the high level of efficiency of the circulatory and respiratory systems, as the training program for the compound physical characteristics contributed to the improvement of these variables. Increasing its ability to contract, so the expansion of the chest increased, and this led to an improvement in vital capacity, rapid exhaled air volume, as well as maximum air volume, and this is consistent with what was indicated by Go Forte et al. Al-Fattah ٢٠٠٠AD indicated that the improvement that occurs in the physiological variables is due to the quality of the training load used according to the physiological trends as well as the training methods used.

(٧٠:٢٦)

The researcher also attributes the significance of the differences to the percentage of lactic acid concentration in the blood after the performance of the experimental group to the program applied to this group and the proper rationing, especially in the inter-rest periods during the time period of the program. Rest periods between each exercise and another and a group and another, as the rest period is important

to extract a large amount of blood accents(١٠:١١٦).

The researcher also indicates that this significant increase in the measurement of lactic acid is due to the fact that the training program for the compound physical characteristics applied to the experimental group led to an improvement in the efficiency of the circulatory and respiratory systems in providing oxygen to delay the accumulation of lactic, as well as the improvement of the work of the enzyme lactan dihydrogenins, which stimulates the reaction of lactic acid to Peruvage, because any disturbances in this enzyme affect this transformation process, as well as the use of active rest after completing free boxing with a colleague through light walking exercises, relaxation exercises, as well as flexibility exercises, which affected the speed of lactic elimination as well as recovery recovery and this is what Confirmed by Ehab Sabri in ٢٠٠٠, citing Merle and Stephen.

This is also in agreement with what Abu El-Ala Abdel-Fattah and Ahmed Nasr El-Din ١٩٩٣m (٢) stated that the applied program led to the availability of oxygen, which led to the oxidation of hydrogen ions and brine and ferric acid to carbon dioxide and water, thus

delaying the accumulation and formation of lactic acid.

The models of this study agree with the results of the studies of Wei & Len ١٩٩٧AD, Jamal Ismail ١٩٩٨AD, Ihab Sabry ٢٠٠٠AD (١٠) quoting from Reborn, Mackinnon, Rozenek, and Burke Pierce that the program Training affects the concentration of lactic acid through an improvement in the enzyme dehydrogenase in the blood, and this contributed to an increase in the ability of boxers to quickly get rid of lactic acid and delay the phenomenon of fatigue and speed recovery recovery

The results of this study are also in agreement with the results of the studies of Sameh Abdel-Raouf ١٩٩٢ (١٨), Cahoon & Bollard ١٩٩٥(٦٨), Abadi Rand et al. ١٩٩٦(٥٩), Ehab Sabry ٢٠٠٠(١٠), Ashraf Mustafa ٢٠٠٣ (٩) Saeed Jamil ٢٠٠٤(٢٢) that the training program leads to an improvement in (maximum oxygen consumption, physical efficiency, anaerobic capacity, anaerobic capacity)

Conclusions and recommendations:

Abstracts:

Within the limits of the objectives, hypotheses and results of the study under study, the researcher reached the following conclusions:

1-The application of the training program by increasing the size of the training load under study led to an improvement in the physical and functional variables of young boxers.

2-The application of the training program by increasing the size of the training load under study led to overcoming the "breaking" pause period in the training session, speeding up the return to the optimal training state "sports formatting" and achieving a better level than the players were before the stopping period.

3-The training program, by increasing the size of the training load, led to the identification of the period necessary for adaptation and overcompensation after the pause period, which is six weeks of training.

Recommendations:

- Based on the results of the study and the resulting conclusions, the researcher recommends the following:

1-The necessity of using the same method of forming training load cycles to design similar programs with an increase in the volume of the training load under the same variables under study "training cycle refraction - the available time range."

2-Benefiting from the use of increasing the size of the training load over the

need for early access to the best sports case "sports format" with the modification of practical applications to match the current study.

3-Not to prolong the cycle of increasing the size of the training load at a week for young women, with a return to the usual training load for a period of two weeks to avoid stress and injury to bring about the required adaptation.

4-Conducting similar studies on samples and other competitions.

Scientific references:

1-Abu Al-Ela Ahmed Abdel-Fattah: (١٩٩٧) Sports Training - Physiological Foundations, first edition, Dar Al-Fikr Al-Arabi, Cairo.

2-Abu El-Ela Ahmed Abdel-Fattah, Ahmed Nasr El-Din: (٢٠٢١) Physiology of Physical Fitness, second edition, Dar Al-Fikr Al-Arabi, Cairo.

3- Egyptian Athletics Federation: (٢٠٢١) Directory of Federation competitions, Cairo.

4- El-Sayed Abdel-Maksoud: (١٩٩٥) Sports Training Theories, Directing and Adjusting the Achievement Level, Training Planning, Training Follow-up, Training Objectives and Sports Competitions, Al-Khansa Library, Cairo.

5-Iman Saad Zaghloul, Iman Nasr: (١٩٩٤) Effect of physical loads of different intensity on the functional efficiency of the respiratory circulatory system among runners ١٠٠m and ٢٠٠m, Journal of Sports Sciences and Arts, Volume VI, Issue III, Faculty of Physical Education for Girls, Helwan University.

6- Bitez. c. Thomson: (١٩٩٣) Introduction to Training Theories, International Association of Athletics Federations - Regional Development Center, Cairo.

7- Hussein Dori Abaza: (٢٠٠٠) The effectiveness of using physical exertion of different intensity on the functional efficiency of the respiratory system for swimmers, Journal of Physical Education Research, Volume ٢٢, Number ٥٤, Faculty of Physical Education for Boys, Zagazig University.

8- Zaki Darwish, Adel Abdel Hafez: (١٩٩٧) Encyclopedia of Athletics, Dar Al Maaref, Cairo.

9-Salah Mohsen Naga, Muhammad Muhammad Zaki, Magdi Muhammad Fahim: (٢٠٠٠) Sports Training, Concepts of Foundations of Principles, Absence, Planning, Language Center of the Age, Tanta.

- 10-Ali Fahmy Al-Baik: (١٩٩٥) Planning Sports Training, Manshayat Al-Maaref, Alexandria.
- 11- Kamal Darwish, Othman Refaat, Muhammad Othman: (١٩٩٢) Physical load and adaptation, an analytical reference study, The Scientific Journal of Physical Education and Sports, Volume Three, Faculty of Physical Education for Boys - Helwan University.
- 12-Muhammad Jamal Hamadeh, Salah Mohsen Naga: (٢٠٠٢) Sports Training, Duties, Methods, Planning, Calendar, Language of the Age Center, Tanta.
- 13- Hisham Ahmed Moheeb: (٢٠٠١) The effect of the direction of training load on the level of some sensory-motor abilities during the stage of direct adaptation, Journal of Theories and Applications, No. ٤٣, Faculty of Physical Education for Boys, Alexandria University.
- 14 - Brown , L . , M. Whitehurst , B.Findley , P. Gilbert , D. Groo , and J. Jimenz (1998) : The effect of repetitions and gender on acceleration range of motion during knee extension an isokinetic device . J. strength Gond – Res – 12 . P. 222 – 255 .
- 15 - Bruin , G. , Kuipers , H.A. Keizer , and G.J. Vander vusse . (1994) : Adaptation and overtraining in horses subject to increasing training loads .J. of Applied physiology 76 : 1908 – 1913 .
- 16 - David Martin and Peter Coe .(1997) : Better training for distance runner .uman Kinetics S٠. U.S.A.
- 17 - Fleck , S.J. (1993) :Cardiovascular responses to strength training .In Strength and power in sport,305-315 .Oxford.U.K. Black well Scientific
- 18 - Fleck , S.J. and W.J. Kraemer. (1997) : Designing resistance exercise programs (2nd . ed .) Champaign , 1L : Human Kinetics .
- 19 - Foster , C. and M. Lehmann. (1997) : Overtraining syndrome . In Runing injuries , ed .GN. Guten , 173 – 188 , philadelphia : Sanders .
- 20 - Fry , A.C., W.J. Kraemer , J.M. lynch , N.T. Trip Lett . and L.P. Koziris . (1994) : Does short- term near maximal intensity machine resistance exercise induce overtraining ? Journal of Applied Sport science Reserch 8 : 188 – 191 .
- 21 - Hooper , S.L. , L.T.Mackinnon,A. Howard,R.D.Gordon and A.W. Bachman . (1995) : Markers for monitoring over training and recovery . Medicine and science in sports and Exercise . 27:106 – 112

- 22 - Jeukendrup , A.E., M.K.C. Hesselink , A.C.S Snyder , H. Kuipers and H.A. Keizer. (1992) : Physiological changes in male competitive cyclists after two weeks of intensified training . International journal of sports Medicine 13: 534-541 .
- 23 - Lehmann , M., E.J. Jakob , U. Gastmann and J.M. Steinacker . (1995) : Unaccustomed high mileage compared to high intensity training related performance and neuromuscular responses in distance runners . European Journal of Applied Physiology 70: 457 – 461 .
- 24 - Lormes , W., J.M. Steinacker and M.Lehmann .(1996) : Short – term overtraining for world championships . International journal for sports medicine . Inpreparation .
- 25 - Rowbottom , D.G., D. Keast and A.R. Morton . (1996) . The emerging role of glutamine as an indicator of exercise stress and overtraining . Sports Medicine 21: 80-97 .
- 26 - Richard , B.K., A.C. Fry and M.L. Toole – 1998 : Overtraining in sport . Human Kinetics Pub . Champaign .